

## INDEX TO VOLUME VII

New names and the final members of new combinations are in **bold face type**

- Abena jamaicensis*, 244  
*Abies grandis*, 121  
*Abronia umbellata*, 64  
*Abrus precatorius*, 340  
*Abutilon hirtum*, 238; *periplocifolium*, 238  
*Acacia riparia*, 178  
*Achyla*, 307; *prolifera*, 307  
*Actaea alba*, 64  
*Adiantum tenerum*, 144  
*Aecidiella Triumfettae*, 253  
*Aecidium*, 66, 86, 170, 171, 253, 254; **abscedens**, 315; *Borreriae*, 315; *Cephalanthi*, 22; *Cestri*, 191; *circumscriptum*, 316; *Cissi*, 316; *Clematidis*, 74; *Clibadii*, 317; *compositarum Eupatorii*, 232; *decoloratum*, 317; *desmium*, 175; *detritum*, 254; *expansum*, 317; **favaceum**, 253; *Hyptidis*, 246-248; *Leonotidis*, 245; *luzoniense*, 254; *Nesaeae*, 86, 89; *passifloricola*, 254; *Phyllanthi*, 254; *pulverulentum*, 315; *Randiae*, 315; *Rivinae*, 235; *roseum*, 250; *solaniphilum*, 255; *Sorbi*, 48; *Tournefortiae*, 254; *Traxini*, 22; *tubulosum*, 255; *tucumanense*, 246; *Uleanum*, 255; *verbenicola*, 22; *Wedeliae*, 318  
*Aeschynomene americana*, 330; *sensitiva*, 330  
*Aesculus glabra*, 65  
*Agaricus*, 95, 97-101; *adirondackensis*, 257; *albo-umbilicatus*, 257; *alveolatus*, 291; *anisarius*, 276; *apertus*, 257; *arvensis*, 100, 102, 152, 226; *auratocephala*, 258; *bicolor*, 105; *brumalis*, 264; *caespitosus*, 281; *campestris*, 96, 100-102, 152; *candicans*, 257; *Candolleanus*, 120; *carnosior*, 259; *catinus*, 258; *cellaris*, 276; *clavipes*, 259; *columbanus*, 259; *compressipes*, 259; *conculus*, 102; *concavus*, 259; *connexus*, 260; *cyathiformis*, 259-261; *dealbatus*, 260; *dicolor*, 261; *diminutivus*, 300; *ectypus*, 106; *ectypoides*, 262; *erubescens*, 262; *facifer*, 281; *flaccidus*, 262; *flavidellus*, 263; *fumosus*, 263; *gallinaceus*, 263; *Gerardianus*, 263; *giganteus*, 264; *gilvus*, 264; *Hoffmani*, 264; *illudens*, 281; *infundibuliformis*, 264, 265; *inversus*, 264; *lacrymabundus*, 116; *leptolomus*, 265; *maculosus*, 265; *marmoreus*, 273; *maximus*, 265; *metachrous*, 261; *mollis*, 268; *monadelphus*, 281; *nebularis*, 268; *nudus*, 105; *odorus*, 276; *pachylus*, 280; *panaeolus*, 106; *peltigerinus*, 267; *personatus*, 105; *phyllophilus*, 268; *pileolarius*, 268; *pinarius*, 268; *pinophilus*, 269; *pithyophilus*, 269; *placomycetes*, 152; *Poculum*, 259; *porphyrellus*, 269; *purus*, 118; *radiozonarius*, 270; *reticeps*, 290, 291; *reticulatus*, 291; *rivulosus*, 270; *Rodmani*, 212; *setisedus*, 270, 271; *Sienna*, 271; *sino-picus*, 271; *socialis*, 272; *sordidus*, 106; *splendens*, 272; *squamulosus*, 274; *subhirtus*, 273; *submarmoreus*, 273; *subzonalis*, 274; *tabescens*, 281; *truncicola*, 276; *trullisatus*, 275; *umbilicatus*, 257; *velutinus*, 116; *vilescens*, 276; *violaceolamelatus*, 120; *violaceus*, 105; *virens*, 276; *viridis*, 276  
*Agaricus reticeps* Mont., Notes on, 290  
*Ageratum conyzoides*, 250  
*Agropyron*, 74, 75; *Smithii*, 73, 78  
*Aklema petiolaris*, 236  
*Albizia Lebbeck*, 146  
*Alchornea latifolia*, 328, 329  
*Aleuria badia*, 90; *pustulata*, 92; *syllvestris*, 93; *vesiculosa*, 91  
*Allium reticulatum*, 66  
*Alpinia antillarum*, 339  
*Ambrosia*, 77; *artemisiaefolia*, 77; *psilostachya*, 76, 77, 88; *trifida*, 77, 88  
*Amelanchier*, 78, 84; *canadensis*, 78, 83, 84, 88, 89; *vulgaris*, 83, 89  
*Amsonia salicifolia*, 84  
*Andropogon*, 71, 72, 88; *brevifolius*, 339  
*Anemia hirsuta*, 326  
*Anemone*, 75; *cylindrica*, 73, 88  
*Apios tuberosa*, 66  
*Apiosporium*, 25  
*Apocynum cannabinum*, 65  
*Aquilegia*, 74, 75; *caerulea*, 73; *canadensis*, 83; *flavescens*, 73, 83

- Arabis, 75, 88  
 Arachis hypogea, 322  
 Argemone mexicana, 334  
 Argomyces **insulanus**, 179; Vernoniae, 179, 180  
 Armillaria, 95, 98; mellea, 102, 131, 132, 152, 281; mellea exannulata, 281  
 Arnica, 77  
 Aronia, 83; arbutifolia, 83  
 Arthur, J. C., Cultures of Uredineae in 1912, 1913, and 1914, 61; Uredinales of Porto Rico based on collections by F. L. Stevens, 168, 227, 315  
 Arthur, J. C., & Fromme, F. D., The taxonomic value of pore characters in the grass and sedge rusts, 28  
 Arundinaria, 22  
 Asclepias, 40, 41; curassavica, 240-242; speciosa, 40, 41; verticillata, 66  
 Ascomycetes, Structural parallelism between spore-forms in the, 21  
 Ascospores of Endothia parasitica, The effect of continued desiccation on the expulsion of, 126  
 Ascotricha, 289  
 Ascrista monticola, 340  
 Ascus-producing species, Penicillium avellaneum, a new, 284  
 Aspergillus, 284  
 Aspidium marginale, 176  
 Aster, 70, 75, 76, 79, 81, 83; Drummondii, 79, 81, 83; ericoides, 72; multiflorus, 72; Novae-Angliae, 72; paniculatus, 65, 69, 70, 75, 76, 79, 81, 83, 84, 87, 88; Tripoli, 70; Tweedyi, 81, 83, 88  
 Auerswaldia, 340; palmicola, 340  
 Axonophus compressus, 326  
 Bagnisiella, 338  
 Baptisia bracteata, 66  
 Basanacantha, 315  
 Bauhinia, 185; divaricata, 185; pauciflora, 184, 185  
 Bidens andicola, 195; leucantha, 195, 196; pilosa, 195, 196; tereticaulis, 195  
 Bixa orellana, 148, 327  
 Bjerkandera adusta, 299  
 Blechnum Brownii, 249  
 Blechnum patula, 320; purpurea, 320  
 Bolbitis varicolor, 214  
 Boletus from California, A new, 44  
 Boletinus, 100; pictus, 300, 305  
 Boletus **californicus**, 215; griseus, 166; ornatus, 166  
 Borreria levis, 249, 250; verticillata, 249, 250  
 Botryosphaeria, 24  
 Boudiera, 199  
 Bouteloua, 31  
 Brodiaea, 63; pauciflora, 85, 89  
 Bromus, 74, 75  
 Brown, H. P., A timber rot accompanying Hymenochaete rubiginosa (Schr.) Lév., 1  
 Buettneria lateralis, 323  
 Caesalpinia, 184  
 Cajan Cajan, 186  
 Cajanus indicus, 186, 187  
 California, A new bolete from, 44  
 Calkins, William Wirt, amateur mycologist, 57  
 Callirhoe digitata, 64, 65, 81, 82; involucreta, 65, 81, 82  
 Calonectria graminicola, 25  
 Canavalia ensiformis, 176  
 Canna, 234, 235; coccinea, 234; glauca, 234; indica, 234; portoricensis, 234  
 Cardiospermum grandiflorum, 236; Halicacabum, 236; microspermum, 236  
 Carduus, 77, Flodmanii, 77, 78  
 Carex, 31, 62, 69, 70, 76, 78, 81, 83, 86; arctata, 67, 87; crinita, 67, 87; extensa, 70; filiformis, 86, 89; Goodenowii, 76; intumescens, 75, 88; maritima, 62; pubescens, 66, 67, 87; retrorsa, 69-87; scoparia, 69, 70, 87; tenuis, 67, 87; tribuloides, 83, 88; vulpinoidea, 79, 81, 87  
 Cassia chamaecrista, 66; quinquangulata, 321  
 Cauloglossum, 100  
 Cayaponia americana, 192, 335; racemosa, 192  
 Cenchrus echinatus, 228, 229, 339; myosuroides, 339; viridis, 228, 229  
 Centella asiatica, 334  
 Cercospora, 41; clavata, 41  
 Ceriomyces affinis, 300; auriporus, 300; communis, 44; crassus, 132, 151; ferruginatus, 152, 300, 306; miniato-olivaceus, 152; nebulosus, 300; retipes, 166; scaber, 152; subglabripes, 300; viscidus, 300, 305  
 Cerotelium Canavaliae, 169, 176  
 Cerrena unicolor, 299  
 Cestrum laurifolium, 191; macrophyllum, 191  
 Chaetochloa brevispica, 231; imberbis, 231; onurus, 231; purpurascens, 231; scandens, 228; setosa, 227, 231; verticillata, 230  
 Chaetomium, 289  
 Chamaecyparis thuyoides, 83, 86, 89

- Chamaesyce hirta*, 190; *hypericifolia*, 190; *prostrata*, 190; *serpens*, 190  
*Chanterel aurantiacus*, 152, 300;  
*Chantarellus*, 152; *infundibuliformis*, 300, 305; *umbonatus*, 300, 305  
 Characters in the grass and sedge rusts, The taxonomic value of pore, 28  
*Chelone glabra*, 64, 65  
*Chlorophyllum Molybdites*, 152  
*Chrysopsis villosus*, 72  
*Cicuta maculata*, 83  
*Cissus sicyoides*, 147, 316  
*Citromyces*, 134  
*Claudopus*, 34-37; *depluens*, 36, 37, 290; **mephiticus**, 290; *nidulans*, 290, 300; **subdepluens**, 36, 37  
*Claudopus*, A new mephitic, 290  
*Claudopus*, A parasitic species of, 34  
*Clavaria*, 305; *cinerea*, 299; *flava*, 151; *fusiformis*, 299; *muscigena*, 49; *pinophila*, 299; *pistillaris*, 299  
*Clematis*, 75; *Douglasii*, 73, 83; *Drummondii*, 82, 83, 88; *ligusticifolia*, 73, 74, 88; *virginiana*, 73, 74, 83  
*Clibadium erosum*, 317; *surinamense*, 317  
*Clitocybe*, 256, 262, 266, 277, 280, 291, 301; *adironackensis*, 257, 258; *aggregata*, 280; *albicastanea*, 278; *albidula*, 257; *albiformis*, 278; *albo-umbilicata*, 257; *ampla*, 263; *angustissima*, 280; *aperta*, 257; *aquatica*, 281; *Arnoldi*, 271; *atrialba*, 278; *avellanealba*, 278; *biformis*, 258, 265; *Broadwayi*, 280; *brumalis*, 261; *brunescens*, 278; *caespitosa*, 258; *candicans*, 257; *candida*, 271, 280; *catina*, 258; *centralis*, 257; *cerussata*, 280; *chrysocephala*, 258; *clavipes*, 259, 266, 278, 301; *columbana*, 259; *compressipes*, 259; *con-cava*, 259; *connexa*, 260; *cuticolor*, 278; *dealbata*, 258, 260, 266, 275, 278; *dealbata sudorifica*, 274; **di-color**, 260, 278; *difformis*, 280; *ditopoda*, 273, 280; *ditopus*, 259; **Earlei**, 261; *eccentrica*, 261, 279, 301; *ectypa*, 280; *ectypoides*, 262, 263; *elixia*, 280; *erubescens*, 262; *fellea*, 262; *flaccida*, 262; *flavidella*, 263; *fragrans*, 280; *fumosa*, 263; *fuscipes*, 263; *gallinacea*, 263; *geotropia*, 265, 280; *Gerardiana*, 263; *gigantea*, 264; *griseifolia*, 278; *Harperi*, 278; **hiemalis**, 264; *hirneola*, 280; *Hoffmani*, 264; *hondensis*, 278; *illudens*, 115, 131, 132, 152, 259, 281; *incrustedata*, 280; *infundibuliformis*, 257, 258, 264, 273, 301; *in-*  
*ornata*, 280; *inversa*, 258, 262, 264, 265, 272, 274, 278; *leptoloma*, 265; *maculosa*, 264, 265; *marginata*, 282; *maxima*, 264, 265; *media*, 266; *megalospora*, 157, 266; *metachroa*, 260, 261; *mexicana*, 280; *microspora*, 279; *monadelphina*, 281; *morbifera*, 260, 266, 275; *multiceps*, 152, 266, 267, 272; *multiformis*, 267; *murinifolia*, 278; *nebularis*, 34, 266, 268; *niveicolor*, 280; *nobilis*, 267; *obobata*, 280; *oculata*, 278; *odora*, 277; *opaca*, 280; *oreades*, 278; *oregonensis*, 278; *parasitica*, 281; *parilis*, 280; *Peckii*, 278; *peltigernia*, 267; *phylophila*, 268; *phylophiloides*, 214, 268; **pileolaria**, 268; *pinariaria*, 268; *pinophila*, 269; *pithophylla*, 269; *porphyrella*, 269; *pruinosa*, 269, 280; *pulcherrima*, 270; *pusilla*, 279; *radi-ozonaria*, 270; *rancidula*, 270; *regularis*, 270; *revoluta*, 282; *rivulosa*, 270, 280; *robusta*, 271, 280; *rubro-tincta*, 280; *setiseda*, 271; *sinopica*, 269, 271, 272, 277, 279, 301; *sino-picoides*, 271; *socialis*, 272, 279; *sphaerospora*, 282; *splendens*, 272; *stipitata*, 278; *subcandicans*, 278; *subconcava*, 271, 272; **subconnexa**, 272; *subcyathiformis*, 273; *subdi-topoda*, 273, 301; *subfumosipes*, 278; *subhirta*, 273; *subinversa*, 278; *subinvoluta*, 280; **submar-morea**, 273; *subnigricans*, 274; *sub-similis*, 273; *subsocialis*, 279; **sub-squamata**, 272, 274; *sudorifica*, 260, 266, 274; *subzonalis*, 274; *sulphu-rea*, 275; **tenebricosa**, 275; *testa-ceoflava*, 280; *Trogii*, 260; *troy-ana*, 280; *trullaeformis*, 280; *trul-lisata*, 275; *truncicola*, 265, 276; *tuba*, 280; *tumulosa*, 280; *variabilis*, 278; *vilescens*, 276; *violaceifolia*, 278; *virens*, 260, 276, 277, 301; *washingtonensis*, 278  
*Clitocybe* in North America, The genus, 256  
*Clitocybe megalospora*, The validity of, 157  
*Clitoria cajanifolia*, 189; *glycinoides*, 189; *rubiginosa*, 189  
*Clusia rosea*, 336  
*Coccoloba*, 145  
*Coccomyces*, 25, 27; *hiemalis*, 45; *lu-tescens*, 45; *prunophorae*, 45  
*Coleosporium*, 84, 169; *Elephantopo-dis*, 171; *Ipomoeae*, 80, 172; *Plu-mierae*, 169, 172; *Solidaginis*, 80; *Vernoniae*, 80, 84, 89  
 Collections by F. L. Stevens, Uredi-

- nales of Porto Rico based on, 168, 227, 315  
*Colletotrichum*, 23, 38, 39, 40, 149;  
*destructivum*, 38; *salmonicolor*, 40; *solanicum*, 39; *Trifolii*, 38  
*Colletotrichum* and *Phoma*, New species of, 38  
*Collomia*, 77; *linearis*, 77, 88  
*Collybia*, 291, 301; *acervata*, 301, 305; *confluens*, 301; *dryophila*, 301, 305; *longipes*, 132; *maculata*, 301, 306; *platyphylla*, 115, 118, 301; *radicata*, 157, 158, 301, 305; *retigera*, 291; *scabriuscula*, 301; *strictipes*, 301; *tuberosa*, 131, 132, 301, 302  
*Colocasia*, 144  
*Coltricia perennis*, 34, 299; *tomentosa*, 299  
*Commelina elegans*, 182, 329; *nudi-flora*, 183; *virginica*, 182, 329, 330  
Conard, Henry S., The structure and development of *Secotium agaricoides*, 94  
Continued desiccation on the expulsion of ascospores of *Endothia parasitica*, The effect of, 126  
*Convolvulus*, 193; *nodiflorus*, 192  
*Coprinus*, 98, 100; *atramentarius*, 34, 152; *comatus*, 152; *finetarius*, 301; *micaceus*, 45, 102, 152, 301  
*Cordia corymbosa*, 337  
*Cordyceps militaris*, 298  
*Coremium*, 134; *silvaticum*, 134  
*Coriolus abietinus*, 299; *versicolor*, 299  
*Corticium coeruleum*, 132  
*Cortinari*, 221, 223, 302; *anomalus*, 223; *armillatus*, 301; *erythrinus*, 223; 302; *lilacinus*, 302, 305; *purpurascens*, 302; *roseipallidus*, 221; *semisanguineus*, 302, 305  
*Cortinellus*, 256; *decorus*, 164, 275  
*Corydalis aurea*, 66  
*Cosmos caudatus*, 195  
*Cracca cinerea*, 177  
*Crataegus cernuus*, 79, 89; *punctatus*, 78; *Pringlei*, 78, 88  
*Craterellus*, 110; *borealis*, 110; *cornucopioides*, 299, 305  
*Crepidotus*, 36, 37  
*Crinipellis*, 156; *alnicola*, 156; *echinulata*, 156; *squamifolia*, 156; *sublivida*, 156; *zonata*, 270  
*Cronartium Quercus*, 80, 89  
*Crotalaria retusa*, 335  
*Croton lucidus*, 147  
*Crucibulum vulgare*, 305  
*Cryptoporus*, 121, 123, 124; *volvatus*, 121, 123, 124, 125, 155  
*Cryptoporus volvatus*, Notes on, 121  
*Cryptosporella*, 24  
*Cudonia lutea*, 298  
Cultures of Uredineae in 1912, 1913, and 1914, 61  
Cup-fungi, Photographs and descriptions of, —  
I. *Peziza*, 90; II. *Sepultaria*, 197  
*Cuphea*, 323; *Parsonsia*, 323  
*Curcubitar*, 27; *Laburni*, 24  
*Cylindrosporium*, 25, 45  
*Cyperus*, 231, 318; *cayennensis*, 231; *laevigatus*, 231; *mutisii*, 231; *polystachus*, 231; *radiatus*, 231; *sphaeclatus*, 231; *surinamensis*, 231  
*Cyphella*, 110  
*Daedalea confragosa*, 299  
*Daldinia concentrica*, 298  
*Dasyscypha*, 9  
*Dasyscypha foveolata*, 241  
*Decodon*, 66; *verticillatus*, 86, 89  
*Delphinium*, 73; *Geyeri*, 73  
*Dendropanax arboreum*, 149  
Descriptions of cup-fungi, Photographs and, —  
I. *Peziza*, 90; II. *Sepultaria*, 197  
Desiccation on the expulsion of ascospores of *Endothia parasitica*, The effect of continued, 126  
*Desmodium axillare*, 188; *Scorpiurus*, 188  
Development of *Secotium agaricoides*, The structure and, 94  
*Dicaeoma*, 28; *canaliculatum*, 231; *ciliata*, 319; *cubense*, 251; *deformatum*, 229; *Eleocharidis*, 232; *Paspali*, 230; *pulcherrimum*, 238; *radicans*, 319  
*Dictyophora duplicata*, 153  
*Dieffenbachia*, 144  
*Dimerina*, 337; *Jacquiniae*, 337  
*Dimerium*, 335, 336, 337; *Cayaponiae*, 335, 340; *grammodes*, 335, 337; *melioloides*, 336, 340; *Stevensii*, 337  
*Dimerosporium*, 335; *lateritium*, 336  
*Dipodia rigida*, 250; *maritima*, 250  
*Diprechidium leve*, 230  
*Dioscorea*, 320; *grandiflora*, 320  
*Ditaca aecia*, 74; *palustris*, 64, 65  
*Doellingeria umbellata*, 65  
*Dolicholus reticulatus*, 186, 187; *texanus*, 186  
*Dolichos*, 187; *Lablab*, 331; *reticulatus*, 187  
*Dorstenia Contrajerva*, 327  
*Dothidea grammodes*, 335; *perisporioides*, 335, 336; *seminata*, 335  
*Dothidella grammodes*, 335  
*Dryopteris patens*, 176, 325; *Poiteana*, 325

- Dulichium arundinaceum*, 81, 87  
Durability of greenheart (*Nectandra Rodiaei* Schomb.), Tests on the, 204
- Edible and poisonous, Fungi, 151  
Effect of continued desiccation on the expulsion of ascospores of *Endothia parasitica*, The, 126  
*Elasmomyces*, 99, 100; *mattioli*anus, 102  
*Eleagnus argentea*, 65, 73  
*Eleocharis*, 232; *cellulosa*, 232; *geniculata*, 232; *interstincta*, 232  
*Elephantopus angustifolius*, 172; *carolinianus*, 84; *mollis*, 171, 172; *scaber*, 172  
*Eleutheranthera ruderalis*, 251, 252  
*Elfvigia Brownii*, 155, 215; *fomentaria*, 299; *megaloma*, 299  
*Elfvigiella*, 110; *fasciata*, 110, 215  
*Elymus*, 74, 75; *canadensis*, 73, 88; *virginicus*, 82, 88  
*Emilia sonchifolia*, 251, 252  
*Endophyllum Rivinae*, 235  
*Endoptychum*, 99, 100  
*Endothia parasitica*, 24, 27, 126, 130  
*Endothia parasitica*, The effect of continued desiccation on the expulsion of ascospores of, 126  
*Entoloma*, 105, 302; *cuspidatum*, 302; *sericeum*, 302; *strictius*, 302  
*Eocronartium typhuloides*, 49  
*Eragrostis tephrosanthus*, 180  
*Erigeron annuus*, 66, 75  
*Eriosporangium Hyptidis*, 247; *tucumanense*, 246  
*Erisyphe*, 46, 328  
*Ernodea littoralis*, 250  
*Erythrina glauca*, 322, 323; *Lithospermae*, 146; *micropteryx*, 146, 323  
*Erythroxylon areolatum*, 320, 321; *Coca*, 321; *havanense*, 320, 321  
*Eugenia buxifolia*, 148; *Jambos*, 239  
*Eupatorium odoratum*, 148; *polyodon*, 250; *villosum*, 251  
*Euphorbia corollata*, 65; *cyparissias*, 65; *hirta*, 190; *petiolaris*, 236  
*Euthamia*, 70; *graminifolia*, 70, 75, 83, 84, 87  
*Exogonium repandum*, 149  
Expulsion of ascospores of *Endothia parasitica*, The effect of continued desiccation on the, 126
- Favolus variegatus*, 289  
Ferax group of the genus *Saprolegnia*, The, 307  
*Ficus*, 174, 175; *Carica*, 174; *glomerata*, 174; *laevigata*, 174  
*Fimbristylis*, 232; *diphylla*, 232; *feruginea*, 232  
Fink, Bruce, William Wirt Calkins, amateur mycologist, 57  
*Fischeria crispiflora*, 242  
*Fistulina*, 155; *hepatica*, 151  
Fitzpatrick, Harry Morton, A parasitic species of *Claudopus*, 34  
*Flammula*, 223; *penetrans*, 302; *polychroa*, 207, 208; *sapinea*, 302; *spumosa*, 302  
Fomes *Abramsianus*, 215; *amarus*, 155; *annosus*, 132, 207, 208; *Brownii*, 215; *everhartii*, 207, 208; *fomentarius*, 207; *igniarius*, 155, 207, 208; *lobatus*, 207, 208; *nigriscanus populinus*, 155; *pinicola*, 207, 208; *putearius*, 13; *robiniae*, 208; *roseus*, 207, 208, 299; *turbinatus*, 215; *ungulatus*, 299  
Fomitiporia *prunicola*, 300  
Fromme, F. D., Arthur, J. C., & The taxonomic value of pore characters in the grass and sedge rusts, 28  
*Fuirena*, 319; *umbellata*, 319  
*Fuligo septica*, 298  
*Fulvifomes*, 110; *calcitratus*, 215; *Cedrelae*, 215; *cinchoensis*, 215; *dependens*, 215; *extensus*, 215; *grenadensis*, 215; *hydrophilus*, 215; *jamaicensis*, 215; *linteus*, 215; *mel-leicinctus*, 215; *pseudosenex*, 215; *sarcitus*, 215; *sublinteus*, 215; *subpectinatus*, 215. *Swieteniae*, 215; *troyanus*, 215; *Underwoodii*, 215; *yucatanensis*, 215  
Fungi edible and poisonous, 151  
Fungi, Illustrations of, — XX, 115; XXI, 163; XXII, 221  
Fungi, Luminescence in the, 131  
Fungi, Preliminary list of Upper St. Regis, 297  
*Fusarium conglutinans*, 49
- Galera Hypnorum*, 302; *tener*, 302  
Garman, Philip, Some Porto Rican parasitic fungi, 333  
Genus *Clitocybe* in North America, The, 256  
Genus *Lepista*, The, 105  
*Gibberella*, 27; *Saubineti*, 25  
*Gloeophyllum hirsutum*, 300  
*Gloeoporus*, 155  
*Gloeosporium*, 23, 40  
*Glomerella*, 23  
*Glycine reticulata*, 186  
*Gnomonia*, 24  
*Gossypium barbadense*, 175; *brasilense*, 175; *hirsutum*, 175  
*Gouania domingensis* 237, 329; *latifolia*, 238; *lupuloides*, 237, 329;

- polygama, 237, 329; tomentosa, 237, 329  
 Grass and sedge rusts, The taxonomic value of pore characters in the, 28  
 Greenheart (Nectandra Rodiaei Schomb.), Tests on the durability of, 204  
 Grifolia Berkeleyi, 46; frondosa, 151  
 Grindelia squarrosa, 72  
 Guarea trichiloides, 149  
 Guettarda ovalifolia, 334  
 Guignardia, 24  
 Guilandina crista, 146  
 Gutierrezia Sarothrae, 72, 88  
 Gymnopilus farinaceus, 222; spumosus, 222  
 Gymnosporangium, 48; Betheli, 78, 88; Botryapites, 83; clavariaeforme, 79, 88; durum, 78, 88; Ellisii, 86, 89; gracilens, 78, 88; Nelsoni, 78, 88; nidus-avis, 83, 89  
 Gyrophila nimbata, 107  
 Gyrophragmium, 100  
 Hamaspora, 87  
 Hapalopilus rutilans, 300  
 Heald, F. D., & Studhalter, R. A., The effect of continued desiccation on the expulsion of the ascospores of Endothia parasitica, 126  
 Hebeloma, 223; palustre, 214  
 Helianthus angustifolius, 65  
 Helicteres jamaicensis, 147  
 Heliomyces, 156  
 Helminthosporium, 109; gramineum, 49; teres, 49  
 Helvella cochleata, 90, 91; Infula, 298  
 Hemidiodia ocimifolia, 250, 315  
 Herbarium, Marking types in the mycological, 108  
 Herpotrichia, 210, 211; nigra, 23, 210, 211  
 Herpotrichia nigra and associated species, Observations on, 210  
 Heteropteryx, 240  
 Heterotrichum cymosum, 337, 338  
 Hibiscus militaris, 64, 82  
 Humphrey, C. J., Tests on the durability of greenheart (Nectandra Rodiaei Schomb.), 204  
 Hyalopora, 170  
 Hydnocystis, 197, 199  
 Hydnum Caput-ursi, 151, 299; ochraceum, 299; repandum, 151; zonatum, 299  
 Hydrophyllum capitatum, 73; Fendleri, 64, 73; virginicum, 64  
 Hygrophorus, 259, 262; chlorophanus, 302; coccineus, 302, 305; miniatus, 302, 305; pratensis, 262  
 Hymenaea, 321; Courbaril, 321  
 Hymenochaete, 9; rubiginosa, 1-4, 13, 14-18  
 Hymenochaete rubiginosa (Schrad.) Lév., A timber rot accompanying, 1  
 Hymenoclea monogyra, 85, 89  
 Hypholoma, 47, 98, 102; appendiculatum, 120, 152; Boughtoni, 117; Candolleum, 120; delineatum, 117; lacrymabundum, 116, 117; perplexum, 152; rigidipes, 117; rugocephalum, 117  
 Hypomyces Ipomoeae, 25  
 Hyptis atrorubens, 246; capitata, 247; latanifolia, 248; pectinata, 246; suaveolens, 246  
 Illustrations of fungi—XX, 115; XXI, 163; XXII, 221  
 Index to American Mycological Literature, 51, 113, 159, 217, 293, 341  
 Indigofera Anil, 177; suffruticosa, 177  
 Inga vera, 177  
 Inocybe, 302; euthelella, 214; infida, 152  
 Inonotus Leei, 155, 215; leprosus, 215; ludovicianus, 110; porrectus, 215; radiatus, 300  
 Ipomoea, 80; acuminata, 243; Batatis, 172; carolina, 243; cathartica, 243; littoralis, 172; Nil, 172; pandurata, 80; stolonifer, 172; trichocarpa, 243; triloba, 243  
 Iresine elatior, 235; paniculata, 235  
 Iris versicolor, 64  
 Ischnoderma fuliginosum, 300  
 Jacquemontia nodiflora, 192, 193, 237; tamnifolia, 172  
 Jacquinia barbasco, 337  
 Jambos Jambos, 239  
 Janipha Manihot, 190  
 Jatropha Curcas, 331; gossypifolia, 331; Manihot, 190  
 Juncus balticus, 76, 88  
 Juniperus scopulorum, 78, 88; sibirica, 79, 89; utahensis, 78, 88; virginiana, 83, 89  
 Klebahnia, 184, 196; Bidentis, 196  
 Koeleria, 75; cristata, 75, 88  
 Kuehneola, 169, 170, 174, 175; Fici, 174; Gossypii, 175  
 Kyllinga caespitosa, 231; pumila, 231  
 Laccaria, 256, 269; laccata, 275, 302, 305; striatula, 302  
 Lachnea hemisphaerica, 299; scutellata, 299  
 Laciniaria Langloisii, 84; punctata, 71, 84; scariosa, 77

- Lactaria*, 302; *camphorata*, 302; *deliciosa*, 152; *Gerardii*, 302; *Hibbardae*, 225; *lactiflua*, 152, 166; *lig-nyota*, 225, 302, 305; *mucida*, 302; *oculata*, 302; *parva*, 302; *piperata*, 152, 225, 302; *plinthogala*, 164; *subdulcis*, 302, 305; *theiogala*, 302; *torminosa*, 302, 306; *turpis*, 301, 302; *varia*, 302; **Volkertii**, 165  
*Lactuca intybacea*, 324  
*Laetiporus speciosus*, 132, 151  
*Lamprospora*, 199  
*Lantana aculeata*, 244; *Camara*, 243, 244, 334; *crocea*, 244; *involucrata*, 244; *odorata*, 244; *stricta*, 244; *trifolia*, 244  
*Lasiacis divaricata*, 180; *Sloanei*, 181; *Swartziana*, 181, 339;  
*Lasiophæra Coulteri*, 210  
*Lentinellus*, 156  
*Lentinula*, 156, 291; **reticeps**, 291  
*Lentinus*, 156; *caespitosus*, 281; *coch-leatus occidentalis*, 216; *lecomtei*, 207, 208; *lepideus*, 119, 207; *umbilicatus*, 214  
**Lentodiellum**, 157, 215, 216; **conca-vum**, 216  
*Lentodium*, 156; *squamosum*, 119  
*Lenzites betulina*, 207, 208; *sepiaria*, 207-209  
*Leonotis nepetaefolia*, 245  
*Leotia lubrica*, 299  
*Lepargyræa canadensis*, 65  
*Lepiota*, 119; *americana*, 152; *amian-thina*, 118, 302; *clypeolaria*, 302, 305; *fuscusquamea*, 302; *meleagris*, 99, 102; *naucina*, 152, 302; *procera*, 152  
*Lepista*, 105; **domestica**, 105, 106; *panacola*, 105, 106; *personata*, 105  
*Lepista*, The genus, 105  
*Leptonia serrulata*, 302  
*Leptostroma*, 23  
*Limacella illinita*, 302  
*Lind's work on the Rostrup Herbarium*, 42  
*List of Upper St. Regis fungi*, Preliminary, 297  
*Lithospermum angustifolium*, 71; *officinale*, 65  
*Lophodermium*, 23  
*Luminescence in the fungi*, 131  
*Lycoperdon*, 100; *atropurpureum*, 305; *cyathiforme*, 152, 305; *gemmaum*, 152, 305; *pulcherrimum*, 305; *pyri-forme*, 305; *separans*, 305; *subincarnatum*, 305; *Wrightii*, 305  
*Lycopus americanus*, 71, 88  
*MacOwenites*, 100  
*Macropodium macropus*, 299  
*Malvastrum*, 82; *coccineum*, 66, 82  
*Man, A new fungus*, *Phialophora verrucosa*, pathogenic for, 200  
*Manihot Manihot*, 190; *utilissima*, 190  
*Manisuris granularis*, 230  
*Marasmiellus*, 156, 157; *inconspicuus*, 156; *juniperinus*, 156  
*Marasmius*, 155, 156, 301; **alnicola**, 156; *campanulatus*, 303; **distanti-folius**, 156; **echinulatus**, 156; **inconspicuus**, 156; **juniperinus**, 156; **niduliformis**, 156; *oreades*, 152; 303; *rotula*, 303; **squamifolius**, 156; **sublividus**, 156; **submulti-ceps**, 156  
*Marasmius in North American Flora*, *Russula* and, 155  
*Marking types in the mycological herbarium*, 108  
*Medlar, E. M., A new fungus*, *Phialophora verrucosa*, pathogenic for man, 200  
*Meibomia*, 189; *adscendens*, 335; *axillaris*, 188; *Scorpiurus*, 188; *tortuosum*, 188, 189  
*Melampsoropsis*, 175  
*Melanoleuca*, 118, 256, 267, 273; *albissima*, 274, 303, 306; *fumidella*, 165; *melaleuca*, 303; *personata*, 152; *rutilans*, 303; *Russula*, 222; *sordida*, 106  
*Melanomma*, 144  
*Melanthera aspera*, 194; *canescens*, 194, 195; *deltoides*, 195; *hastata cubensis*, 195  
*Melasmia*, 23  
*Melothria guadalupensis*, 192  
*Menyanthes*, 66  
*Mephitic Claudopus, A new*, 290  
*Merulius*, 155; *lacrymans*, 11, 206-209; *tremellosus*, 207, 208  
*Mesosphaerum atrorubens*, 246, 247; *capitatum*, 247; *latanifolium*, 248; *pectinatum*, 246, 247; *spicatum*, 246; *suaveolens*, 246, 247  
*Metastelma lineare*, 242; *palustre*, 243; *parviflorum*, 242; *Schlechtendahl*, 242  
*Miconia*, 338; *prasina*, 337; *Sintensii*, 337, 338; *laevigata*, 333, 337  
*Microcera*, 23  
*Mikania*, 317; *cordifolia*, 317  
*Milesia*, 176; **columbiensis**, 175; **con-similis**, 176; **Kriegeriana**, 176  
*Milesina columbiensis*, 175; *Kriegeriana*, 176  
*Mimosa ceratonia*, 183, 184  
*Mitracarpus portoricensis*, 250  
*Mitrula vitellina*, 299  
*Momisia iguanaea*, 145  
*Monadelphus*, 256, 277, 281; *caespi-*



- tosus, 281; illudens, 258, 281, 282;  
**marginatus**, 282; **revolutus**, 282;  
**sphaerosporus**, 282  
 Montagnites, 100  
 Morchella esculenta, 151  
 Muhlenbergia, 82, 181; Porteri, 81  
 Murrill, W. A., Fungi edible and  
 poisonous, 151; The genus Clitocybe  
 in North America, 256; The ge-  
 nus Lepista, 105; Illustrations of  
 Fungi—XX, 115, XXI, 163, XXII,  
 221; Luminescence in the fungi,  
 131; Marking types in the myco-  
 logical herbarium, 108; A new bo-  
 lete from California, 44; Prelimi-  
 nary list of Upper St. Regis fungi,  
 297  
 Mutinus, 100  
 Mycena, 280, 303; Leaiana, 303; pura,  
 117, 269, 303  
 Mycological herbarium, Marking types  
 in the, 108  
 Mycologist, amateur, William Wirt  
 Calkins, 57  
 Myrica cerifera, 87, 89  
 Mytilidion, 210, 211; fusisporum, 211  
  
 Napaea, 82; dioica, 82;  
 Nectria discophora, 25; galligena, 25;  
 megalospora, 336  
 Neopectia Coulteri, 210, 211  
 Nephrolepis rivularis, 175  
 Neurocarpum cajanifolium, 189  
 Neurolaena lobata, 252  
 New ascus-producing species, Pencil-  
 lium avellaneum, A, 284  
 New Bolete from California, A, 44  
 New fungus, Phialophora verrucosa,  
 pathogenic for man, A, 200  
 New mephitic Claudopus, A, 290  
 News and Notes, 109, 212, 288  
 and Reviews, 45, 155  
 New species of Colletotrichum and  
 Phoma, 38  
 Nigredo, 28, 183, 192, 194; appendi-  
 culata, 185; **columbiana**, 194; Com-  
 melinae, 182; Dolicholi, 186; Erag-  
 rostidis, 180; Hedyasari-paniculati,  
 188, 189; Helleriana, 192; leptoder-  
 ma, 180; major, 181; Neuro-  
 carpi, 189; proëminens, 190; Rhyn-  
 cosporae, 182; Scleriae, 182  
 North American Flora, Russula and  
 Marasmius in, 155  
 North America, The genus Clitocybe  
 in, 256  
 Notes on Agaricus reticeps Mont.,  
 290  
 Notes on Cryptoporus volvatus, 121  
 Nyctalis, 34  
  
 Observations on Herpotrichia nigra  
 and associated species, 210  
 Octospora pustulata, 92  
 Oenothera grandiflora, 46; Tracyi, 46  
 O'Gara, P. J., New species of Colletot-  
 richum and Phoma, 38  
 Olyra latifolia, 229  
 Omphalia 263, 304; campanella, 303;  
 chrysophylla, 303; fibula, 303; um-  
 bellifera, 304  
 Onagra biennis, 75  
 Onosmodium occidentale, 73  
 Ophionectria, 27; coccicola, 23, 25  
 Orton, C. R., Structural parallelism  
 between spore-forms in the Asco-  
 mycetes, 21  
 Otidea, 91  
  
 Panaeolus solidipes, 163  
 Pandanus, 150  
 Panellus, 156; flabellatus, 156; jala-  
 pensis, 156; subcantharelloides, 156  
 Panicum, 339; barbinode, 181; cap-  
 illare, 65; divaricatum, 180; lana-  
 tum, 181; maximum, 144; molle,  
 181; ovalifolium, 229; Sloanei, 181;  
 trichoides, 229; virgatum, 65  
 Panus concavus, 216; **flabellatus**, 156;  
 incandescens, 282; **jalapensis**, 156;  
 meruliceps, 291; stypticus, 131,  
 132; **subcantharelloides**, 156  
 Parallelism between spore-forms in  
 the Ascomycetes, Structural, 21  
 Parasitic fungi, Some Porto Rican,  
 333  
 Parasitic fungi—I, Studies in Porto  
 Rican, 143  
 Parasitic species of Claudopus, A, 34  
 Parodiella, 335, 337; melioides, 336  
 Paspalum, 230, 339; compressum,  
 326; conjugatum, 319; fimbriatum,  
 230; Helleri, 230, 326; Humboldtianum,  
 319, 326; orbiculatum, 230;  
 paniculatum, 230, 326; pilosum,  
 230; plicatum, 230, 326; Schre-  
 berianum, 230; Underwoodii, 339  
 Passiflora rubra, 254  
 Pathogenic for man, A new fungus,  
 Phialophora verrucosa, 200  
 Paxillus, 105; atrotomentosus, 304;  
 involutus, 152, 304, 305; Lepista,  
 105; panuoides, 304  
 Peltigera, 267  
 Pelvetia, 213  
 Penicillium, 134, 135, 136, 138, 284;  
 africanum, 135, 142; **avellaneum**,  
 284; brevicaulis, 134; candidum,  
 284; claviforme, 134; duclauxi, 134,  
 140, 141; expansum, 135; funicu-  
 losum, 142; luteum, 135, 136, 137,  
 140, 284; pinophilum, 135, 136, 140,



- 142; *purpureogenum*, 135, 136, 140, 142; *purpureogenum* var. *rubri-sclerotium*, 137; *rugulosum*, 140, 141, 142; *silvaticum*, 134
- Penicillium avellaneum*, a new ascus-producing species, 284
- Penicillium luteum-purpureogenum* group, The, 134
- Peridermium*, 79, 84; *carneum*, 80, 84; *Cerebrum*, 80, 214; *fusiforme*, 62, 79, 80, 84, 89;
- Petitia domingensis*, 333
- Peziza*, 90, 197; *abietina*, 299; *arenicola*, 197; *assimilata*, 92; *badia*, 90, 91, 93; *bufonia*, 93; *cochleata*, 90, 91; *heterophylla*, 73; *pustulata*, 92, 93; *sepulta*, 197, 198; *sylvestris*, 93; *tanacetifolia*, 64; *umbrina*, 92; *vesiculosa*, 91, 93
- Phaeolus sistotremoides*, 300
- Phakopsora*, 173, 175; *Phyllanthi*, 332; *Vitis*, 173
- Phallus*, 100
- Phaseolus*, 186, 187; *adenanthus*, 185; *lunatus*, 335; *prostratus*, 187; *vulgaris*, 185
- Phialophora**, 202; *verrucosa*, 202, 203
- Phialophora verrucosa*, pathogenic for man, A new fungus, 200
- Philadelphus coronarius*, 78, 88; *Ke-teleerii*, 78, 88
- Philibertella clausa*, 243
- Philibertia*, 243
- Pholiota*, 304; *candicans*, 152; *Johnsoniana*, 226; *praecox*, 214; *squarrosa*, 304
- Phoma*, 41, 143; *rostrata*, 41
- Phoma*, New species of *Colletotrichum* and, 38
- Photographs and descriptions of cup-fungi—  
I. *Peziza*, 90; II. *Sepultaria*, 197
- Phragmidium*, 175
- Phragmites*, 22
- Phyllachora*, 337, 338; *Andropogonis*, 339, 340; *dalbergicola perforans*, 340; *graminis*, 338, 339, 340; *nitens*, 339; *paulensis*, 340; *perforans*, 340; *peribebuensis*, 337, 338, 340; *Renalmiae*, 339; *sphaerosperma*, 339
- Phyllanthus distichus*, 332; *grandifolius*, 332; *nobilis*, 254
- Phyllosticta*, 143; *adianticola*, 144; *araliana*, 148; *bixina*, 148; *boringensis*, 147; *cissicola*, 147; *Colocasiae*, 144; *colocasicola*, 144; *commelinicola*, 144; *divergens*, 146; *Erythrinae*, 146; *erythrinicola*, 146; *Eugeniae*, 148; *eupatoriicola*, 148; *glaucispora*, 149; *guanicensis*, 146; *Guareae*, 149; *Ipomoeae*, 149; *Lagenariae*, 149; *momisiana*, 145; *nericola*, 149; *Nerii*, 149; *pandanicola*, 150; *Panici*, 144; *Pithecolobii*, 145; *Pithecolobii monensis*, 145; *portoricensis*, 147; *Sacchari*, 144; *Sechii*, 149; *Stevensii*, 147
- Physalis pubescens*, 66
- Physopella*, 170, 173, 175, 330; *Aechynomenis*, 330; *Fici*, 174; *ficina*, 174; *Vitis*, 173
- Picea*, 210; *Engelmanni*, 210
- Pieters*, A. J., The *ferax* group of the genus *Saprolegnia*, 307
- Pinus*, 169; *palustris*, 80, 89; *taeda*, 79, 80, 84, 89
- Piptoporus suberosus*, 300
- Pithecolobium Unguis-cati*, 145
- Pityrogramma calomelanos*, 325, 334
- Pleospora*, 25, 27, 49
- Pleuropus abortivus*, 152
- Pleurotopsis*, 156; *niduliformis*, 156
- Pleurotus*, 36, 37, 291; *caespitosus*, 281; *cubensis*, 156; *lampas*, 282; *noctiluens*, 282; *olearius*, 282; *orizabensis*, 156; *sapidus*, 152; *subbarbatulus*, 156
- Plicaria badia*, 90; *pustulata*, 92
- Plowrightia*, 25
- Pluchea odorata*, 324; *purpurascens*, 324, 325
- Plumiera alba*, 173; *Krugii*, 172; *obtusata*, 172; *rubra*, 173
- Pluteus alveolatus*, 291; *cervinus*, 152, 304
- Poinsettia heterophylla*, 190
- Poisonous, Fungi edible and, 151
- Polymarasmus*, 156, 157; *submulti-ceps*, 156
- Polyplocium*, 100
- Polyporus*, 34, 35; *adustus*, 207; *arcularius*, 214; *caudicinus*, 132; *elegrans*, 300; *graminicola*, 215; *Leei*, 215; *Marbleae*, 215; *McMurphyi*, 155; *obtusatus*, 207, 208; *perennis*, 34-37; *porrectus*, 215; *resinosus*, 207, 208; *Schweinitzii*, 214; *sulphureus*, 207, 208; *volvatus*, 125; *Zelleri*, 155
- Polystictus hirsutus*, 207, 208, 209; *prolificans*, 208; *versicolor*, 207, 208
- Pore characters in the grass and sedge rusts, The taxonomic value of, 28
- Porodaedalea Pini*, 111, 132, 300
- Porothelium*, 155
- Porto Rican parasitic fungi, Some, 333

- Porto Rican parasitic fungi—I, Studies in, 143
- Porto Rico based on collections by F. L. Stevens, Uredinales of, 168, 227, 315
- Preliminary list of Upper St. Regis Fungi, 297
- Prosopodium appendiculatum, 178
- Psalliotia, 101
- Pseudopeziza, 24
- Pseudotsuga taxifolia, 121
- Psidium, 240; Guajava, 239
- Psychotria patens, 323
- Pteris, 334
- Puccinia, 22, 28, 62, 67, 76, 170, 245, 322, 323; Agropyri, 73, 74, 82, 88; albiperidia, 67, 68, 69, 78, 87; alternans, 74; Andropogonis, 72; angustata, 64, 70, 71, 88; angustatoidea, 62; appendiculata, 178; Arechavelatae, 236; Beckii, 179; Blechi, 249; Boutelouae, 30; **Cameliae**, 227; Campulosi, 29; canaliculata, 231, 318; Cannae, 233; Caricis-Asteris, 32, 69, 76, 87; Caricis-Solidaginis, 69, 79, 87; Caricis-Strictae, 32; Cenchr. 32, 228; Chaetochloae, 230; compacta, 240, 241; **concrescens**, 240, 241; Convolvuli, 192; crassipes, 243; Cryptandri, 64; Cynanchi, 242; Cyperi, 231; deformata, 229; Dolichii, 186; Dulichii, 66, 84, 81, 87; Eleocharidis, 232; Eleutherantherae, 251; Ellisiana, 71, 72, 88; emaculata, 65; Emiliae, 251; epiphylla, 29, 32; esclavensis, 32; Euphorbiae, 236; extensicola, 70, 76, 79, 80, 81, 87; Fimbristylidis, 232; Fuirenae, 319; Gonolobi, 243; Gouaniae, 237, 329; gouaniicola, 238; graminis, 62; gregaria, 241; Grossulariae, 66, 67, 69, 76, 87; hemisphaerica, 324; heterospora, 238; Huberi, 229; Hyptidis, 247, 248; **insititia**, 248; Ipomoeae, 243; inflata, 236; Jambosae, 239, 240; karelica, 32; Latanae, 243; lateritia, 249; leonotidicola, 245; Leonotidis, 245; leptospora, 29; levis, 230; macropoda, 235; McClatchiana, 64; medellinensis, 246, 247, 249; Melicae, 30; minutissima, 86, 89; monoica, 75, 88; Muhlenbergiae, 81, 82, 88; Nesaee, 86; nodosa, 85, 89; obliqua, 242, 243; oblitterata, 74; Pammeli, 65; Panici, 65; panicola, 180; paradoxica, 29; Paspali, 230; peridermiospora, 22; poculiformis, 32; Psidii, 239, 240; quadripurula, 76, 87; Raunkiaerii, 235; Ribesii-Caricis, 69; Rivinae, 235; Rompelii, 240; rosea, 250; rubigo-vera, 62; salviicola, 249; sanguinolenta, 240; **scleritica**, 232; Seymouriana, 22, 30; solida, 251, 252; sphaerospora, 242, 243; splendens, 85, 89; Sporoboli, 31, 32; Stipae, 72, 88; substriata, 230; Synedrellae, 251; tageticola, 251; Thaliae, 233; Thwaitessii, 238; tomipara, 74; tosta, 64, 66, 81, 82, 84, 88; triarculata, 30; Tridacis, 251; **tucumanensis**, 247; uniporula, 67, 68, 69, 78; Urbaniana, 244; Urticae, 32; Vernoniae, 179; Vernoniae-mollis, 179; versicolor, 30; vexans, 31, 32; Vilfae, 22; vulpinoides, 62, 63, 66, 79, 80, 84, 87; xanthopoda, 233
- Pucciniastrum, 169, 175
- Puccinosira, 253; pallidula, 170, 253;
- Pustularia vesiculosa, 91
- Pyropolyporus Abramsianus, 155, 215; cinnabarinus, 300; conchatus, 300; igniarius, 111, 300; igniarius nigricans, 111
- Quamoclit coccinea, 172
- Quercus Phellos, 80, 89; rubra, 79, 80, 89
- Rajania, 320; cordata, 320
- Randia, 315; aculeata, 315
- Ravenelia, 178, 184, 322, 329; Caesalpiniae, 183; capituliformis, 328; caulicola, 176; Indigoferae, 177; Ingae, 177; **Stevensii**, 178
- Resupinatus, 156; cubensis, 156; orizabensis, 156; subbarbatulus, 156
- Reticularia Lycoperdon, 298
- Rheosporangium aphanidermatum, 288
- Rhizina inflata, 213
- Rhizoctonia, 39, 48; solani, 48
- Rhodopaxillus, 105; panaculus, 107; personatus, 105; sordidus, 106
- Rhynchospora, 62; aurea, 182; cyperoides, 182; distans, 182; micrantha, 182; polyphylla, 182
- Rhyncosia, 187; longeracemosa, 187; reticulata, 186; Senna, 187; texana, 186
- Rhytisma, 23
- Ribes, 69, 76; Cynosbati, 66, 78, 87;
- Rivina humilis, 235; octandra, 235
- Romell, Lars, Lind's work on the Rostrop Herbarium, 42
- Rosellinia melioloides, 336
- Rostkovites **californicus**, 44, 215; granulatus, 152, 300, 305; subaureus, 300, 305

- Rostrup Herbarium, Lind's work on the, 42
- Rostrupia Scleriae, 233
- Rot accompanying Hymenochaete rubiginosa (Schräd.) Lév., A timber, 1
- Ruellia strepens, 65
- Russula, 155, 156, 166, 221, 222, 225, 273, 304; albida, 304; bifida, 224; compacta, 304, 305; crustosa, 224; depallens, 304; emetica, 152, 304, 306; foetens, 152, 304, 306; furcata, 224; lutea, 304, 305; Mariae, 152; pectinatoides, 167; virescens, 152, 224
- Russula and Marasmius in North American Flora, 155
- Rusts, The taxonomic value of pore characters in the grass and sedge, 28
- Rytilix granularis, 230
- Sabicea aspera, 324
- Saccharum, 144
- Salvia occidentalis, 249
- Saprolegnia, 307; anisospora, 308; declina, 308, 312; dioica, 308; ferax, 307-314; hypogyna, 308; mixta, 307-314; monoica, 307-309, 312-314; Thureti, 307, 308; torulosa, 307, 308
- Saprolegnia, The ferax group of the genus, 307
- Sarcosphaera, 199
- Schizophyllum commune, 208
- Schizophyllum, 156
- Schlegelia brachyantha, 339
- Schroeteriaster, 170
- Scirpus atrovirens, 70, 88; Eriophorum, 71; fluviatilis, 83, 88; microcarpus, 64, 71; rubroinctus, 64, 71; sylvaticus, 64, 71
- Scleria, 182, 233; Baldwinia, 233; hirtella, 233; pauciflora, 233; pterota, 182; setacea, 233; verticillata, 233
- Scleroderma, 167; aurantium, 153, 305; tenerum, 167
- Sclerotinia, 24
- Scopulariopsis, 134
- Scutiger griseus, 300; hispidellus, 155
- Scytinotus, 156; distantifolius, 156
- Seaver, Fred J., Observations on Herpotrichia nigra and associated species, 210; Photographs and descriptions of cup-fungi—I. Peziza, 90; II. Sepultaria, 197
- Secium edule, 149
- Secotium, 98, 99-103; agaricoides, 94, 97, 99-102; coarotatum, 100; erythrocephalum, 99; Gueinzii, 100; Mattirolanus, 99; melanosporum, 100; olbium, 100, 103; Warnei, 94
- Secotium agaricoides, The structure and development of, 94
- Sedge rusts, The taxonomic value of pore characters in the grass and, 28
- Senecio Douglasii, 72; lugens, 72; obovatus, 81; spartioides, 72, 88
- Septoria, 333; aquilina, 334; asiatica, 334; Chelidonii, 334; Guettardae, 334, 340; Lantanae, 334, 340; Miconiae, 333, 340; Petisiae, 333, 340; Pityrogrammae, 334; Verbenae, 334
- Sepultaria, 197, 198, 199; arenicola, 198; Longii, 199
- Setaria scandens, 228; setosa, 227; verticillata, 230
- Sida, 239; glutinosa, 238; humilis, 238, 239; nervosa, 238; procumbens, 238; spinosa, 238; urens, 238, 239
- Sidalcea, 82
- Silphium perfoliatum, 77
- Sitodrepa panicea, 124
- Sium cicutae-folium, 83, 88
- Solanum torvum, 255; tuberosum, 39
- Solenia, 155
- Solidago, 63, 75, 76, 79, 81, 83; canadensis, 65, 70, 72, 75, 79, 81, 83, 84, 87, 88; glaberrima, 79, 87; missouriensis, 79; mollis, 72, 79, 87; rugosa, 79, 83, 87
- Some Porto Rican parasitic fungi, 333
- Sparganophorum Vaillantii, 325
- Spartina, 22; Michauxiana, 77, 88
- Spathularia velutipes, 299
- Species of Claudopus, A parasitic, 34
- Species of Colletotrichum and Phoma, New, 38
- Spermocoe aspera, 250; tenuior, 250
- Sphaeralcea, 82; incana, 64, 81, 88; lobata, 64, 65, 82, 88
- Sphaeria canaliculata, 231; melioides, 336; perisporioides, 335, 336
- Spongipellis borealis, 300; fragilis, 110
- Spore-forms in the Ascomycetes, Structural parallelism between, 21
- Sporobolus, 82, 181; asperifolius, 81, 88; cryptandrus, 64; diandrus, 181; indicus, 181; longifolius, 66
- Stachytarpheta jamaicensis, 244; strigosa, 244
- Steironema ciliatum, 77; lanceolatum, 77
- Stemmodontia trilobata, 318
- Stenolobium Stans, 178
- Stereum, 2; fasciatum, 207, 208, 209; frustulosum, 13-17; gausapatum,

- 207, 208; hirsutum, 16; rameale, 207, 208
- Stigmaphyllon lingulatum*, 236; *periplocifolium*, 236; *Sagraeanum*, 236
- Stipa comata*, 72, 88
- Strobilomyces strobilaceus*, 152
- Stropharia*, 95, 98; *ambigua*, 103; *bi-lamellata*, 226; *coprinophila*, 34; *semiglobata*, 304
- Struchium Sparganophorum*, 325
- Structural parallelism between spore-forms in the Ascomycetes, 21
- Structure and development of *Secotium agaricoides*, The, 94
- Strumella*, 46; *coryneoides*, 46
- Studhalter, R. A., Heald, F. D., & The effect of continued desiccation on the expulsion of the ascospores of *Endothia parasitica*, 126
- Studies in Porto Rican parasitic fungi—I, 143
- Suillelus luridus*, 152, 300, 306
- Synedrella nodiflora*, 251
- Tagetes patula*, 251
- Taxonomic value of pore characters in the grass and sedge rusts, The, 28
- Tecoma Stans*, 178
- Tectella*, 156
- Teramnus uncinatus*, 331
- Tests on the durability of greenheart (*Nectandra Rodiaei* Schomb.), 204
- Tetrazygia*, 337, 338
- Thalia*, 234, 235; *geniculata*, 234
- Thalictrum*, 75; *alpinum*, 73; *dioicum*, 74, 83; *Fendleri*, 64, 73, 83
- Telephora laciniata*, 299; *perdix*, 13
- Thom, Charles, The *Penicillium luteum-purpureum* group, 134
- Thom, Charles, & Turesson, G. W., *Penicillium avellaneum*, a new ascus-producing species, 284
- Timber rot accompanying *Hymenochaete rubiginosa* (Schr.) Lév., A., 1
- Tournefortia hirsutissima*, 254
- Tradescantia multiflora*, 183
- Trametes Pini*, 13; *abietis*, 13, 14, 17; *Pini*, 214; *robinophila*, 207
- Tremella lutescens*, 299; *mycetophila*, 299
- Tremellodon gelatinosum*, 299
- Tricholoma*, 105, 256, 291; *cellare*, 277; *multipunctum*, 165; *nudum*, 105; *panaeolum*, 105, 107; *personnatum*, 105, 270; *rancidulum*, 270; *Russula*, 222; *Sienna*, 277; *sordidum*, 601
- Tridax procumbens*, 252
- Trifolium carolinianum*, 85, 89; *hybrum*, 38; *pratense*, 38, 39
- Triopteris lingulata*, 236
- Trisetum*, 75; *subspicatum*, 75, 88
- Triumfetta*, 253; *grandiflora*, 253; *Lappula*, 253; *rhomboidea*, 253; *semitriloba*, 147, 253
- Tsuga heterophylla*, 121
- Turesson, G. W., Thom, Charles, & *Penicillium avellaneum*, a new ascus-producing species, 284
- Tylopilus felleus*, 151, 300, 306
- Types in the mycological herbarium, Marking, 108
- Typhula muscicola*, 49
- Tyromyces caesius*, 300; *chioneus*, 300; *graminicola*, 215; *guttulatus*, 300; *lacteus*, 300
- Upper St. Regis fungi, Preliminary list of, 297
- Urechites lutea*, 149
- Uredinales of Porto Rico based on collections by F. L. Stevens, 168, 227, 315
- Uredineae in 1912, 1913, and 1914, Cultures of, 61
- Uredo, 170, 330; *Aeschynomenis*, 330; *Agerati*, 250; *amaniensis*, 195; *Arachidis*, 322; *balaensis*, 249; *basipora*, 245; *bidenticola*, 195; *Bidentis*, 195; *biocellata*, 324; *Bixae*, 327; *Cabreriana*, 322; *Cajani*, 187; *Cameliae*, 227, 228; *cancerina*, 246; *Cannae*, 233; *capituliformis*, 328; *Cephalanthi*, 324; *Cestri*, 191; *Chaetochloae*, 230; *Commelinaceae*, 182; *Commelinae*, 182; *Commelyneae*, 329; *concors*, 330; *Cupheae*, 323; *Desmodii-tortuosi*, 189; *Diehromenae*, 319; *Dioscoreae*, 320; *Dolichi*, 186; *Erythroxylonis*, 320; *Eugeniarum*, 239, 240; *fallaciosa*, 323; *fenestrata*, 170, 332; *Fici*, 174; *ficeicola*, 174; *ficina*, 174; *flavidula*, 239; *Fuirenae*, 319; *gemmata*, 193, 237; *Gossypii*, 175; *Gouaniae*, 329; *Gymnogrammes*, 170, 325; *Hymenaeae*, 321; *ignobilis*, 181; *Jacquemontiae*, 193; *Janiphae*, 190; *jatrophicola*, 331; *Kyllingiae*, 231; *leonotica*, 245, 246; *Leonotidis*, 245; *lutea*, 321; *morica*, 174; *Myrtearum*, 239, 240; *neurophylla*, 240; *nigropunctata*, 320; *nootkatensis*, 48; *pamparum*, 187; *paspalica*, 319, 326; *Phyllanthi*, 332; *proximella*, 324; *rubescens*, 327; *sabiceicola*, 323; *Sparganophori*, 325; *Stevensiana*, 326; *striolata*, 235; *subneurophylla*,

- 240; *superficialis*, 326; *Teramni*, 331; *vicina*, 325; *Violae*, 173; *Vitis*, 173
- Uromyces*, 22, 28, 31, 32, 170, 184, 240, 322, 323; *acuminatus*, 77, 88; *appendiculatus*, 185, 187; *Arachidis*, 322; *Bidentis*, 195, 196; **Caesalpiniae**, 183; *Cestri*, 191; *columbianus*, 194; *Commelinae*, 182, 330; **densus**, 196; *dichrous*, 190; *Dolicholi*, 186, 187; *elegans*, 85, 89; *Eragrostidis*, 180; *Euphorbiae*, 189; *euphorbiicola*, 190; *gemmatus*, 192, 237; *Hedysari-paniculati*, 188; *Hellerianus*, 192; **ignobilis**, 181; *insularis*, 189; *jamaicensis*, 184; **Janiphae**, 190; *Jatrophae*, 191; *Junci*, 76, 88; *leptodermus*, 180; *major*, 181; *malvacearum*, 238; *malvicola*, 238; *Neurocarpi*, 189; *pedatatus*, 72; *perigynius*, 75, 76, 83, 88; *pianhyensis*, 227, 325; *picatus*, 238; *Poae*, 32; *proëminens*, 189; *pulcherrimus*, 238; *quadripodula*, 76; *Rhynchosporae*, 65, 182; *Scirpi*, 83, 88; *Scleriae*, 182, 233; *Sidae*, 238; *solidus*, 188; *Sporoboli*, 66; *Thwaitesii*, 238; *uniporulus*, 32, 67, 68
- Vaginata agglutinata*, 152; *plumbea*, 152, 304, 305; *plumbea strangulata*, 305
- Vagnera stellata*, 66
- Valerianodes jamaicensis*, 244; *strigosa*, 244
- Validity of *Clitocybe megalospora*, The, 157
- Value of pore characters in the grass and sedge rusts, The taxonomic, 28
- Venenarius Caesareus*, 152; *cothurnatus*, 152; *phalloides*, 152, 305, 306; *Frostianus*, 305; *muscarius*, 152, 305, 306; *rubens*, 152; *solitarius*, 152; *spretus*, 152
- Venturia*, 25, 26, 27
- Vermicularia*, 39
- Vernonia*, 80, 84; *albicaulis*, 179; *boringuensis*, 180; *fasciculata*, 80, 84, 89; *longifolia*, 179; *phyllostachya*, 180
- Vigna*, 186, 187; *repens*, 185; *vexillata*, 185
- Vincetoxicum*, 243
- Viola*, 71, 72; *cucullata*, 71, 88; *Nuttallii*, 71, 88; *premulaefolia*, 71; *striata*, 72
- Vitis*, 173; *vinifera*, 173
- Volvaria Loveiana*, 34
- Wallrothiella Arceuthobii*, 289
- Wedelia*, 227, 325; *carnosa*, 318; *lancoolata*, 325; *reticulata*, 227; *trilobata*, 318
- Wissadula periplocifolia*, 238, 239
- Work on the Rostrup Herbarium, Lind's 42
- Xanthium*, 318; *canadense*, 77
- Xylaria Hypoxylon*, 131, 132; *polymorpha*, 299
- Xylopia*, 241
- Young, Esther, Studies in Porto Rican parasitic fungi, 143
- Zanthoxylum americanum*, 66
- Zeller, Sanford M., Notes on *Cryptoporus volvatus*, 121